

Reducing surgical site infections in Caesarean Sections

we protect lives
worldwide



INTRODUCTION


42% of babies in the UK are delivered by caesarean section [NHS Digital, 2024].

One of the most frequently recorded complications of caesarean section is Surgical Site Infection (SSI); yet they have been described as "the most preventable HCAI" [Leaper, 2015].

A number of measures including skin and wound care can be effective in helping prevent infection.

CAESAREAN SECTION

Caesarean section is one of the most frequently performed surgical procedures [Stanirowski, 2016] and rates are increasing. In England in 1980, 9% of deliveries were by C-section, [Wloch, 2020] compared to 42% in 2023 [NHS digital, 2024].



Caesarean section is usually recommended where a vaginal delivery (VD) can pose risks to either the mother or baby, or both, including:


- Prolonged or obstructed labour
- Foetal distress
- Elevated blood pressure
- Raised glucose levels
- Abnormal presentation/position of the baby [Angolile, 2023].

There has been a recent increase in planned 'Maternal Request Caesarean Sections' (MRCS), and NICE Guidance on Caesarean Birth (2024) recommends that women who request an MRCS should be able to give birth this way, if appropriate.



SURGICAL SITE INFECTIONS

Surgical site infection (SSI) describes wound infections following invasive surgical procedures, usually within 30 days of surgery [Haque, 2018].




They are one of the most common healthcare adverse events, [Haque, 2018].

SSI:

- Accounts for 14.5% of all hospital-acquired infections in the UK [Health Innovation, W England, 2023]
- Leads to an estimated 34-226% increase in associated health costs, [Health Innovation, W England, 2023].
- Can be prevented in around 60% of cases, [Collins, 2019].

CAESAREAN SECTION SURGICAL SITE INFECTIONS

Surgical Site Infection (SSI) is one of the most frequent complications following caesarean section with an estimated incidence of around 16% [Health Innovation, W England, 2023]. This is high compared to 8.6% for large bowel surgery and 9.7% for cholecystectomy, [UKHSA, 2022].



Given the short length of hospital stay, the overall rate of SSI following caesarean section could be significantly higher, as 91% of SSIs occur after discharge from hospital, [Public Health, Wales Report, 2018].

As well as causing anxiety and pain for the mother, post-caesarean SSI may extend the period of hospitalisation by 4 days, and cost an additional £3,173 per patient, [Stanirowski, 2016].

A BMI over 30 kg/m² [Wloch, 2012; Eritty, 2022] and diabetes – gestational or pre-existing Type 1 diabetes [Wloch, 2012] are recognised risk factors for SSI.



POST-OPERATIVE WOUND CARE

NICE: Caesarean Birth (2024) offers advice on post-operative wound care, which includes daily wound cleansing and monitoring for signs of infection.

Due to the improved formulation of existing antiseptics with fewer side effects, wound antiseptics is increasingly being considered as an essential factor in the prevention, treatment and management of wound infection, [Barrigah-Benissan, 2022].

A study of 543 women undergoing elective or emergency caesarean section included the use of wound irrigation with octenidine solution as part of an infection prevention strategy, [Stanirowski, 2016].

Following abdominal surgery, a study of 45 patients showed significant improvements in scar appearance, pain, skin elasticity, and incidence of scar hypertrophy following management with octenidine-based hydrogel compared to conventional dry wound dressings, [Matiasek, 2018].

PREVENTING CAESAREAN SECTION SURGICAL SITE INFECTIONS



DECONTAMINATION

Staphylococcus aureus is a known risk factor for developing SSI and has been identified as the causative organism in over 40% of post-caesarean wound infections, [Wloch, 2012].

Up to 30% of people in the UK carry *Staphylococcus*, [Jeans, 2018] and NICE Guidelines [2019] suggest the use of a body wash "before procedures in which *Staphylococcus aureus* is a likely cause of a surgical site infection" as it "may reduce SSIs". Evidence suggests "any additional costs incurred ... are likely to be more than recouped by savings associated with a lower incidence of surgical site infections." [NICE, 2019].

Research supports the efficacy of preoperative decontamination with an octenidine-based antimicrobial. A large scale trial reported a three-fold reduction in the MSSA infection rate, after a 5 day decontamination regime which included octenisan® wash lotion, [Jeans, 2018].



PERIWOUND CARE

The skin surrounding a wound (the periwound area) is particularly vulnerable and although it may appear healthy, periwound complications frequently occur, [Bianchi, 2012].

Periwound damage can contribute to delayed wound healing and increased wound size, [Woo, 2017]. Best practice guidelines recommend cleaning the periwound area thoroughly but gently, using a skin-friendly cleanser (pH v4–6), with disposable moistened cloths or cleansing pads, [LeBlanc, 2021].

POST – DISCHARGE WOUND CARE

Many women are discharged 24 hours after a caesarean delivery [NHS UK, 2024], but most wound infections present in the community.

As part of periwound care, octenidine containing wash mitts could have a role, on intact skin, [Dhoonmoon, 2024]. A 12 month community evaluation found the use of these mitts led to a reduction in the prescription of antibiotics for wound infections, a reduction in unwarranted infections and improvements in the quality of care delivered, [Dhoonmoon, 2020].

SKIN PREPARATION PRIOR TO CAESAREAN SECTION SURGERY

The skin is a major source of pathogens causing SSI and pre-surgery skin preparation with antiseptic agents has been shown to reduce the risk of infection, [Al Maqbali, 2013].

NICE (2024) recommends the use of an alcohol-based chlorhexidine skin preparation before caesarean birth to reduce the risk of wound infections.



CASE STUDIES

CASE STUDY 1

Expecting her first child, 32 year old Zoe was diagnosed as clinically obese with a BMI of 38, and therefore at increased risk of pregnancy complications. She was referred for Consultant-led care and closely monitored for pregnancy-related diabetes and hypertension. Zoe was advised to have an elective caesarean section at 40 weeks. Before admission, she followed a 5-day body-washing regime using octenisan® wash mitts. Following delivery of a baby boy, she was discharged home two days after surgery and continued using wash mitts as part of her periwound care. She found them easy to use, particularly in the skin folds around her caesarean wound site. Zoe and her baby did well and she is already following a gradual weight loss programme.



CASE STUDY 3

Aisha, 28, had experienced difficulties with the vaginal delivery of her first two children. Whilst pregnant with her third child she requested a planned caesarean section, which her Trust agreed to. She used octenisan® body wash for 5 days before being admitted at 39 weeks. The caesarean section delivery of a baby boy was carried out under epidural anaesthetic. 24 hours later, Aisha returned home to the care of the community midwife. She used octenisan® wash mitts daily to keep the periwound skin clean.

Her scar healed quickly and Aisha regularly applied octenicare® repair creme to support skin regeneration and help reduce scarring.



CASE STUDY 2

Natasha went into spontaneous labour at 41 weeks with her second child. On admission, her baby was found to be in the breech position. Despite strong contractions, her labour progressed very slowly. Close monitoring indicated foetal distress; and the baby's heart rate fell below 100bpm. An emergency caesarean section was carried out within 30 minutes and a healthy baby girl was delivered. Daily woundcare included octenilin® irrigation solution and octenilin® wound gel. This regime continued after discharge and no post caesarean complications were recorded.



OCTENIDINE

Octenidine is a broad spectrum antimicrobial that is less susceptible to bacterial resistance [Spencer, 2013]; is effective in a short contact time at low concentrations, [Assadian, 2016]; and is the only antiseptic that remains active locally for up to 48 hours, [Malanovic 2020].

THE OCTENIDINE-CONTAINING RANGE

octenilin® wound irrigation solution removes wound coating and biofilm; and creates an ideal environment for wound healing.

octenilin® wound gel effectively moisturises and cleans wounds to support the natural healing process.

octenisan® wash lotion for mild and gentle whole body washing against MDRO.

octenisan® wash mitts are ready-to-use for cleaning and care of the skin; also for MDRO decontamination.




octenicare® repair creme contains panthenol to support the natural regeneration of the skin and is suitable for epithelialised wounds.

References

- Al Maqbali MA. Preoperative antiseptic skin preparations and reducing SSI. *Br J Nurs*. 2013;22(21):1227-1233.
- Angolillo CM, Max BL, Mushemba J, Mashauri HL. Global increased cesarean section rates and public health implications: A call to action. *Health Sci Rep*. 2023;6(5):e1274. Published 2023 May 18.
- Assadian O. Octenidine dihydrochloride: chemical characteristics and antimicrobial properties. *J Wound Care*. 2016;25(3 Suppl):S3-S6. doi:10.12968/jowc.2016.25.Sup3.S3.
- Barrigah-Benissan K, Ory J, Sotto A, Salipante F, Lavigne JP, Loubet P. Antiseptic Agents for Chronic Wounds: A Systematic Review. *Antibiotics (Basel)*. 2022;11(3):350. Published 2022 Mar 6.
- Bianchi J (2012) Protecting the integrity of the periwound skin. *Wounds Essentials* 7(1): 58–64.
- Chadwick P, et al, Product Evaluation: Case studies: Octenilin Wound Irrigation Solution and Octenilin Wound Gel in practice *Wounds UK, EWMA SPECIAL*, 2016.
- Collins CR, Wick EC. Reflections on the Complexity of Surgical Site Infection Prevention and Detection from an Organizational Lens. *Surg Infect (Larchmt)*. 2019;20(7):577-580.
- Dhooonmoon L, Dyer M. Improving leg ulcer care in the community. *Journal of Community Nursing*, vol. 34, no. 6, Dec. 2020, pp. 40.
- Dhooonmoon L. Necrotising fasciitis: is follow-up supported self-care an option? *Journ Comm Nurs*. 2024; Vol 38, No 1, 24-28.
- Eisenbeiss W, et al. Prospective, double-blinded, randomised controlled trial assessing the effect of an Octenidine-based hydrogel on bacterial colonisation and epithelialization of skin graft wounds in burn patients. *Int J Burns Trauma*. 2012;2(2):71-79.
- Erritty M, Hale J, Thomas J, et al. Evaluation of independent risk factors associated with surgical site infections from caesarean section. *Arch Gynecol Obstet*. 2023;308(6):1775-1783.
- Haque M, Sartelli M, McKimm J, Abu Bakar M. Health care-associated infections - an overview. *Infect Drug Resist*. 2018;11:2321-2333. Published 2018 Nov 15.
- Health Innovation, West of England, Preventing Caesarean Birth Surgical Site Infection, 2023 <https://www.healthinnowest.net/our-work/transforming-services-and-systems/preventing-surgical-site-infections/reducing-surgical-site-infection-caesarean-birth/> Accessed 9th February 2024.
- Jeanes E, Holleyman R, Tate D, Reed M, Malviya A. Methicillin sensitive staphylococcus aureus screening and decolonisation in elective hip and knee arthroplasty. *J Infect*. 2018;77(5):405-409. doi:10.1016/j.jinf.2018.05.012.
- Leaper D. An overview of surgical site infection Review. *Wounds UK, EWMA SPECIAL*, 2015 p 14-19.
- LeBlanc K, Beeckman D, Campbell K et al (2021) Best practice recommendations for prevention and management of periwound skin complications. *Wounds International*.
- Malanovic N, Ōn A, Pabst G, Zellner A, Lohner K. Octenidine: Novel insights into the detailed killing mechanism of Gram-negative bacteria at a cellular and molecular level. *Int J Antimicrob Agents*. 2020;56(5):106146.
- Matiasek J, Kienzl P, Unger LW, Grill C, Koller R, Turk BR. An intraindividual surgical wound comparison shows that octenidine-based hydrogel wound dressing ameliorates scar appearance following abdominoplasty. *Int Wound J*. 2018;15:914–920.
- NHS Digital, 2023, <https://digital.nhs.uk/data-and-information/publications/statistical/maternity-services-monthly-statistics/december-2023-experimental-statistics> Accessed 22nd March 2024.
- NHS Caesarean Section, 2024. <https://www.nhs.uk/conditions/caesarean-section/#> Accessed 22nd March 2024.
- NICE guideline: Caesarean birth. Published: 31 March 2021. Last updated: 30 January 2024. www.nice.org.uk/guidance/ng192 accessed 27th February 2024.
- Public Health Wales NHS Trust. Annual All Wales report on Caesarean Section Surgical Site Infection 2018, Wales.
- Spencer C, Orr D, Hallam S, Tillmanns E. Daily bathing with octenidine on an intensive care unit is associated with a lower carriage rate of methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect*. 2013;83(2):156-159.
- Stanirowski PJ, Bizoń M, Cendrowski K, Sawicki W. Randomized Controlled Trial Evaluating Dialkylcarbamoyl Chloride Impregnated Dressings for the Prevention of Surgical Site Infections in Adult Women Undergoing Caesarean Section. *Surg Infect (Larchmt)*. 2016;17(4):427-435.
- UK Health Security Agency, Surveillance of surgical site infections in NHS hospitals in England April 2021 to March 2022. Published December 2022, Prepared by: Louise Thorn, Theresa Lamagni, Pauline Harrington, Catherine Wloch, Russell Hope and Colin Brown.
- Wloch C, Wilson J, Lamagni T, Harrington P, Charlett A, Sheridan E. Risk factors for surgical site infection following caesarean section in England: results from a multicentre cohort study. *BJOG* 2012;119:1324–1333.
- Wloch C, Van Hoek AJ, Green N, et al Cost-benefit analysis of surveillance for surgical site infection following caesarean section *BMJ Open* 2020;10:e036919.
- Woo KY, Beeckman D, Chakravarthy D (2017) Management of moisture-associated skin damage: A scoping review. *Adv Skin Wound Care* 30(11): 494-501.

Note

The terms mother and mothers have been used throughout this brochure, as this is the way that the majority of those who are pregnant and having a baby will identify. It also includes people whose gender identity does not correspond with their birth sex or who may have a non-binary identity.

-  www.youtube.com/schuelkeChannel
-  www.facebook.com/myschulke
-  [X.com/schulkeUK](https://x.com/schulkeUK)

Schülke & Mayr UK Ltd.
Sheffield S9 1AT | United Kingdom
Phone +44 114 254 35 00
Fax +44 114 254 35 01
www.schuelke.com

**we protect lives
worldwide**